

REPORT DOCUMENTATION PAGE

AFRL-SR-BL-TR-00-

Public reporting burden for this collection of information is estimated to average 1 hour per response, including gathering and maintaining the data needed, and completing and reviewing the collection of information, including suggestions for reducing this burden, to Washington Headquarters, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Project, Washington, DC 20503.

data sources, aspect of this 215 Jefferson 3503.

0436

1. AGENCY USE ONLY (Leave blank)		2. REPORT DATE		3. REPORT DATE 1 January 1995 - 30 June 1998	
4. TITLE AND SUBTITLE Environment Conditions Responsible for Solar Activity				5. FUNDING NUMBERS F49620-95-1-0008	
6. AUTHOR(S) Prof Peter A. Sturrock					
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Stanford University, Varian Center for Space Science and Astrophysics Stanford, CA 94305-4060				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES) AFOSR 801 N. Randolph Street, Room 732 Arlington, VA 22203-1977				10. SPONSORING/MONITORING AGENCY REPORT NUMBER F49620-95-1-0008	
11. SUPPLEMENTARY NOTES					
12a. DISTRIBUTION AVAILABILITY STATEMENT Approved for Public Release				12b. DISTRIBUTION CODE	
13. ABSTRACT (Maximum 200 words) During the seven years that this program was active at Stanford University, the group working with Professor Sturrock investigated many aspects of the conditions responsible for solar activity. Their results are presented in detail in their publications. A list of publications, sorted by topic, is included in the report.					
14. SUBJECT TERMS "Magnetic Field Structure & Analysis"; "Coronal Structure & Heating"; "Flares, CMEs, and Eruptions"; "Prokinences", "Solar Internal Dynamics"				15. NUMBER OF PAGES	
				16. PRICE CODE	
17. SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT		

20000925 085

DTIC QUALITY INSPECTED 4

Standard Form 298 (Rev. 2-89) (EG)
Prescribed by ANSI Std. Z39.18
Designed using Perform Pro, WHS/DIOR, Oct 94

Environmental Conditions Responsible for Solar Activity

Final Technical Report

For the Period January 1, 1991 through June 30, 1998

Principal Investigator: Professor Peter A. Sturrock

Grant No.: AF-F49620-95-1-0008

Program Manager: Dr. Henry R. Radoski

STAFF

P.I. Professor Peter A. Sturrock

Other Faculty: Guenther Walther

Senior Research Associates: Spiro Antiochos, Taeil Bai and George Roumeliotis

Research Associates: James Klimchuk and Michael Wheatland

Post-Graduate Students: Slava Glukhov and Lisa Porter

Summary

During the seven years that this program was active at Stanford University, the group working with Professor Sturrock investigated many aspects of the conditions responsible for solar activity. Their results are presented in detail in their publications. A list of publications, sorted by topic, is included in this report. The following is a very brief summary of the major advances.

Magnetic Field Structure and Analysis

One of the major problems concerning solar activity and its effect on Earth is that of coronal mass ejections (CMEs). These are believed arise from the sudden expansion of magnetic-field configurations. The final state appears to

be that of an open magnetic field. Sturrock proved analytically that the open state is in fact maximum-energy state of a semi-infinite magnetic field configuration. This result has the important consequence that it is not possible for magnetic field to be driven into a completely open state purely by magnetic stresses. This has proved to be an important guide in subsequent thinking about CMEs.

Glukhov, Klimchuk, Porter and Roumeliotis further developed the magneto-frictional method (created at Stanford in 1986) for calculating magnetic-field configurations. This technique was applied to study of the evolution of magnetic field configurations as they were progressively stressed. We examined a case related to CMEs, using a spherical geometry and an initially dipolar magnetic field, and then examined the development of the magnetic field when the northern and southern hemispheres are rotated in opposite directions, to simulate the foot-point shearing that is believed to be responsible for some CMEs. We obtained the surprising result that steady and slow motion of the footpoints could lead to a sudden eruption of the magnetic field. This may explain the origin of a certain class of CMEs.

Since our investigations led us to calculate the structure of highly stressed force-free field configurations, we decided to examine the asymptotic forms of force-free fields of translational and cylindrical symmetry. Sturrock derived formulas showing how the dimensions and energy of the field vary with continued surface displacements. Glukhov and Porter found that the analytical formulas obtained in this way are a good approximation to the results of our computer calculations.

Towards the end of this program, Roumeliotis developed what we consider to be an important new idea for reconstructing force-free magnetic fields from photospheric vector magnetograph data. This is called the “optimization function” approach. Subsequent study of this method has shown that it could lead to a practical procedure for calculating the coronal magnetic-field structure from photospheric vector magnetograph data. This could be very

helpful since many models of coronal magnetic fields have been investigated, to see if they lead to flares or CMEs. Hence one will be able to take advantage of those studies if one can calculate the actual structure of the coronal magnetic field, based on observational data.

Coronal Structure and Heating

There is an extensive literature concerning coronal heating, which nevertheless remains a major problem in solar physics. Since the corona is hotter and denser in active regions (which have strong magnetic fields) than in quiet regions (which have weak magnetic fields), there is a strong case that the corona is heated by MHD (magnetohydrodynamic) waves. However, there has been the difficulty that fast-mode MHD waves, which can carry sufficient energy to heat the corona, do not dissipate in the corona. Porter and Klimchuk were able to show that it is indeed possible for the corona to be heated by MHD waves, but they must be of much higher frequency than previously believed. This led us to look for processes which might excite high-frequency waves in the corona.

This led Sturrock to develop a new concept for coronal heating, that of "sudden magnetic relaxation." The basic idea is that, if two flux tubes reconnect at or near their foot-points, this will lead to the formation of a large flux tube that is far from equilibrium. The sudden relaxation of that flux tube towards its equilibrium state will then excite sound waves (and probably other types of waves also). These sound waves will then dissipate in the corona and lead to coronal heating.

We have carried out a number of calculations to evaluate this concept. We have developed the idea theoretically, and Parnell made a comparison of the coronal parameters and the relevant photospheric magnetic field strength. There appears to be good agreement between the theory and observational

data, and we find that this concept gives a good explanation both of the energy budget of the corona and of the mass flow into and out of the corona.

Flares, CMEs and Eruptions

Our study of the destabilization of force-free magnetic field configurations led to a promising interpretation of eruptive flares, in terms of the sudden expansion of a slowly stressed force-free field configuration. Sturrock further developed the idea that “tether-cutting” may play an important role in driving prominence eruptions and eruptive flares. The term refers to an analogy: if a hot-air balloon is held down by a number of weak ropes, and if one of the ropes snaps or is cut, the others may be too weak to hold the balloon in place so that they also snap and the balloon lifts off. A prominence may involve a cylindrical flux tube that is held in place by many magnetic connections to the photosphere. If some of these connections break, this may lead to a progressive breaking of the remaining connections so that the prominence lifts off, causing a CME.

Wheatland further developed the avalanche model of solar flares, originally developed by Russ Hamilton and Ed Lu at Stanford. The model involves a large array of cells, and rules according to which sudden energy release in one cell may trigger one or more energy-release events in neighboring cells. Earlier studies of this process implicitly considered only a single active region. Our development took into account the fact that flares occur in many different active regions of different sizes and energies. This led to a more realistic basis for the avalanche model, which should be helpful in future development of this theory.

Prominences

Antiochos and Klimchuk developed a theory for the condensation of coronal gas that can lead to the formation of solar prominences. Glukhov investigated the tearing instability of a neutral current sheet, such as exists

between the oppositely directed flux tubes in a helmet streamer, and developed a self-consistent model for the thermodynamic state of a helmet streamer.

Internal Dynamics

Solar activity is normally regarded as having its origin in events on the solar surface. However, the fact that sunspots develop as part of the overall solar cycle, and the fact that there are other cycles in solar activity, clearly indicates that there are major processes in the solar interior that determine the occurrence and properties of solar activity. We therefore attempted to better understand the nature of periodicity in solar activity, and to find other ways to probe the solar interior. This led Bai and Sturrock to study oscillations that occur with periods of 154 days (the Rieger oscillation), 77 days, and 51 days. We also undertook to search for such periodicities in solar neutrino data, since this is the only way of probing the internal magnetic field of the Sun.

Analysis by Sturrock, Walther and Wheatland led us to the conclusion that the solar neutrino flux does not vary with the solar cycle, contradicting a number of articles that had previously been published. We believe those articles were based on faulty statistics. However, we found evidence that the solar neutrino flux does vary with the rotational period of the Sun. This result, if it is confirmed by later studies, will provide strong evidence that neutrinos have non-zero magnetic moment and that they change their flavor in propagating through magnetized regions in the solar interior. This could potentially provide a method for probing the solar magnetic field deep in the convection zone. If this proves possible, it should lead to advance notice of the development of the solar cycle and it could possibly provide an early warning of the development of new active regions.

PUBLICATIONS
Supported in part by
AF-F49620-95-1-0008

MAGNETIC FIELD STRUCTURE AND ANALYSIS

Sturrock, P.A., 1991, Maximum Energy of Semi-Infinite Magnetic Field Configurations, Ap. J., 380, 655 - 659

Klimchuk, J. A., Canfield, R. C., and Rhoads, J. E., 1992, The Practical Application of the Magnetic Virial Theorem, Ap. J., 385, 327-343

Porter, L. J., Klimchuk, J. A., and Sturrock, P. A., 1992, Cylindrically Symmetric Force-Free Magnetic Fields, Ap. J., 385, 738-745

Roumeliotis, G. , 1993, A new class of exact, nonlinear solutions to the Grad-Shafranov equation, Ap. J., 404, 781 - 787

Roumeliotis, G., and Moore, R.L., 1993, A linear solution for magnetic reconnection driven by converging or diverging flows, Ap. J., 416, 386 - 391

Roumeliotis, G., 1994, A method for reconstructing the coronal magnetic field from vector magnetograph measurements at the photosphere, Solar Active Region Evolution: Comparing Models with Observations, ASP Conf. Series, 68 (eds. K.S. Balasubramaniam and G.W. Simon), 240 - 243

Roumeliotis, G., Sturrock, P.A., and Antiochos, S.K., 1994, A numerical study of the sudden eruption of sheared magnetic fields, Ap. J. 423, 847 - 863

Sturrock, P.A., Antiochos, S.K., Klimchuk, J.A., and Roumeliotis, G., 1994, Asymptotic forms for the energy of force-free magnetic-field configurations of translational symmetry, Ap. J. 431, 870 - 872

Sturrock, P.A., Klimchuk, J.A., Roumeliotis, G., and Antiochos, S.K., , 1994, The asymptotic behavior of force-free magnetic-field configurations, Solar Active Region Evolution: Comparing Models with Observations, ASP Conf. Series, 68 (eds. K.S. Balasubramaniam and G.W. Simon), 219-224

Roumeliotis, G., 1996, The "stress-and-relax" method for reconstructing the coronal magnetic field from vector magnetograph data, Ap J. 473, 1095

Sturrock, P.A., Klimchuk, J.A., & Roumeliotis, G., 1995, Asymptotic analysis of force-free magnetic fields of cylindrical symmetry, Ap. J., 443, 804 - 809

Roumeliotis, G., & Wheatland, M.S., 1997, An optimization approach to reconstructing force-free fields. I. Theoretical basis, BAAS, 29, 920

Wheatland, M.S., & Roumeliotis, G., 1997, An optimization approach to reconstructing force-free fields. II. Numerical results, BAAS, 29, 920

CORONAL STRUCTURE AND HEATING

Roumeliotis, G., 1991, Joule heating as an explanation for the differential emission measure structure and systematic redshifts in the Sun's lower transition region, Ap. J., 379, 392-400

Porter, L.J., Klimchuk, J.A., and Sturrock, P.A., 1994, The possible role of MHD waves in heating the solar corona, Ap. J. 435, 482 - 501

Porter, L.J., Klimchuk, J.A., and Sturrock, P.A., 1994, The possible role of high-frequency waves in heating solar coronal loops, Ap. J. 435, 502 - 514

Klimchuk, J.A., and Porter, L.J., 1995, The Scaling of Heating Rates in Solar Coronal Loops, Nature, 377, 131

Porter, L.J., and Klimchuk, J.A., 1995, Soft X-ray loops and coronal heating, Ap. J. 454, 499

Sturrock, P.A., and Acton, L.W., 1995, Interpretation of SXT data concerning the diffuse corona, BAAS, 27, 964

Sturrock, P.A., Wheatland, M.S., and Acton, L.W., 1995, Interpretation of SXT data concerning the diffuse corona, Proc. IAU Symp. No. 153, Magnetodynamic Phenomena in the Solar Atmosphere (eds. Y. Uchida, T. Kosugi & H.S. Hudson; Dordrecht, Kluwer), 417-418

Wheatland, M.S., and Sturrock, P.A., 1995, Coronal hard X-ray sources in solar flares, BAAS, 27, 991 - 992

Sturrock, P.A., 1996, Coronal Heating by Sudden Magnetic Relaxation, Magnetic Reconnection in the Solar Atmosphere, ASP Conference Series, 111 (eds. R.D. Bentley and J.T. Mariska), 105-111

Sturrock, P.A., 1996, Perspectives on Solar Flares, Solar Drivers of Interplanetary and Terrestrial Disturbances (A.S.P. Conf. Series No. 95, eds. K.S. Balasubramaniam, S.L. Keil & R.N. Smart), 42 - 54

Sturrock, P.A., Wheatland, M.S., and Acton, L.W., 1996, Yohkoh Soft X-Ray Telescope images of the diffuse solar corona, Ap. J. (Letters), 461, L115 - L117

Sturrock, P.A., Wheatland, M.S., and Acton, L.W., 1996, Interpretation of SXT data concerning the diffuse corona, Magnetodynamic Phenomena in the Solar Atmosphere (eds. Y. Uchida, T. Kosugi & H.S. Hudson; Dordrecht, Kluwer), 417-418

Wheatland, M.S., and Sturrock, P.A., 1996, Coronal heating in the quiet corona, BAAS, 28, 869

Wheatland, M.S., Sturrock, P.A. and Acton, L.W., 1997, Coronal heating and the vertical temperature structure of the quiet corona, *Ap. J.* 482, 510

FLARES, AND ERUPTIONS

Moore, R.L., and Roumeliotis, G., 1992, Triggering of eruptive flares: destabilization of the preflare magnetic field configuration, *Eruptive Solar Flares* (eds. Z. Svestka, B.V. Jackson and M.E. Machado; New York: Springer), p. 69

Sturrock, P.A., 1992, The Emerging Picture of Eruptive Solar Flares, *Comments in Astrophysics*, 16, 71 - 85

Sturrock, P.A., 1992, The Emerging Picture of Eruptive Solar Flares, *Proc. IAU Coll. 133* (Iguazu, August 1991) on Eruptive Solar Flares (Springer-Verlag, 1992), 397 – 409

Moore, R., Porter, J., Roumeliotis, G., Tsuneta, S., Shimizu, T., Sturrock, P., and Acton, L., 1994, Observations of enhanced coronal heating in sheared magnetic fields, *Proc. Symp. New Look at the Sun* (Kofu, September 1993). Nobeyama Radio Observatory Report No. 360, 89 – 92

Porter, J., Moore, R. T., Roumeliotis, G., Shimizu, T., Tsuneta, S., Sturrock, P. A., and Acton, L. W., 1994, Microflaring at the feet of large active region loops., *Proceedings of Kofu Symposium* (eds. Shinzo Enome and Tadashi Hirayama), Nobeyama Radio Observatory Report No. 360, 65 – 69

Roumeliotis, G., 1995, A numerical study of the tether cutting model for eruptive flares, *BAAS*, 27, 980

Wheatland, P.A., and Sturrock, P.A., 1996, Avalanche models of solar flares and the distribution of active regions, *Ap. J.* 471, 1044

Wheatland, M.S., & Glukhov, S., 1998, Flare frequency distributions based on a master equation, *Ap. J.* 494, 858

Wheatland, M.S., Sturrock, P.A., & McTiernan, J., 1998, The waiting-time distribution of solar flare hard X-ray bursts, *Ap. J.*, 509, 448

PROMINENCES

Antiochos, S. K., and Klimchuk, J. A., 1991, A New Model For the Formation of Solar Prominences, *Ap. J.*, 378, 372-377

Roumeliotis, G., and Emslie, A. G., 1991, Magnetohydrodynamics of an impulsively heated, hard X-ray emitting filament, *Ap.J.*, 377, 685-693

Glukhov, S., 1996, Tearing instability of a neutral current sheet in the solar corona, *Ap. J.* 469, 936-943

Glukhov, S., 1997, Thermodynamics and morphology of a helmet streamer observed by the Yohkoh soft X-ray telescope, *Ap. J.* 476, 385-391

INTERNAL DYNAMICS

Bai, T., and Sturrock, P.A., 1991, The 154-Day and Related Periodicities as Subharmonics of a Fundamental Period, *Nature*, 350, 141 – 143

Bai, T., 1992, Methods of Periodicity Analysis: Relationship between the Rayleigh Analysis and a Maximum Likelihood Method, *Ap. J.* 397, 584

Bai, T., 1992, The 77-Day Periodicity in the Flare Occurrence Rate of Cycle 22, *Ap. J. (Letters)* 388, L69

Sturrock, P.A., and Bai, T., 1992, Search for Evidence of a Clock related to the Solar 154-day Complex of Periodicities, *Ap. J.*, 397, 337 – 346

Bai, T., 1993, Variability of the Occurrence Frequency as a Function of Hard X-Ray Peak Rate, *Ap. J.*, 404, 805

Bai, T., and Sturrock, P.A., 1993, Evidence for a fundamental period of the Sun and its relation to the 154-day complex of periodicities, *Ap. J.* 409, 476 – 486

Bai, T., 1994, The 51-day periodicity in Cycle 22, *Solar Phys.* 150, 385- 388

Sturrock, P.A., Walther, G. and Wheatland, M.S., 1997, Search for periodicities in the Homestake solar neutrino data, *Ap. J.*, 491, 409 - 413

Walther, G., 1997, Absence of Correlation between the Solar Neutrino Flux and the Sunspot Number, *Phys. Rev. Lett.* 79, 4522-4524.

Sturrock, P.A., 1998, Is the solar cycle due to a dynamo process?, *Proc. SOLER22 Workshop, National Solar Observatory, June 17 - 21, 1996* (eds. J.M. Pap, C. Frohlich & R.K. Ulrich; Kluwer), p. 427

Sturrock, P.A., Walther, G., & Wheatland, M.S., 1998, Apparent latitudinal modulation of the solar neutrino flux, *Ap. J.* 507, 978 – 983